



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\text{Power applied} = \frac{Fs}{t} = \frac{941 \times 300}{24} = 11,762 \text{ ft. lbs. per sec.} = 21 \text{ H.P. (approx.).}$$

283. Proposed by C. N. SCHMALL, New York City.


The maximum length of a certain chain which can be suspended from one end without breaking is l . It is desired to form a catenary with a length $2l/k$ of the chain, the points of support being a distance d apart, in the same horizontal line. Show that the maximum value of d is

$$\frac{2l}{k} (k^2 - 1)^{1/2} \log \left(\frac{k+1}{k-1} \right)^{1/2}.$$

SOLUTION BY A. M. HARDING, University of Arkansas.

Let w = weight per unit length. Then wl = maximum tension the chain will stand.

The tension at the point of support is given by $T \sin \theta = ws$ where s = one-half the length of the chain and θ is the angle that the tangent to the catenary at that point makes with the x -axis.



If we put $T = wl$ and $s = l/k$ we find $\sin \theta = 1/k$. But $\tan \theta = \frac{1}{2}(e^{d/2c} - e^{-(d/2c)})$ where c is the distance along the Y -axis from the origin to the catenary.

Hence we have

$$\frac{1}{2}(e^{d/2c} - e^{-(d/2c)}) = \frac{1}{\sqrt{k^2 - 1}}.$$

From this equation we obtain

$$d = 2c \log \left(\frac{k+1}{k-1} \right)^{\frac{1}{2}}.$$

We have also the intrinsic equation of the catenary $s = c \tan \theta$, from which we obtain

$$c = \frac{l}{k} \sqrt{k^2 - 1}.$$

Whence

$$d = \frac{2l}{k} (k^2 - 1)^{\frac{1}{2}} \log \left(\frac{k+1}{k-1} \right)^{\frac{1}{2}}.$$

Also solved by J. W. COLSON.

NUMBER THEORY.

189. Proposed by V. M. SPUNAR, Chicago, Illinois.

If p and $p_1 = 2^p - 1$ are primes, then are the numbers of the sequence $p_1 = 2^p - 1$, $p_2 = 2^{p_1} - 1$, $p_3 = 2^{p_2} - 1$, \dots , $p_n = 2^{p_{n-1}} - 1$ all primes?

REMARKS BY R. D. CARMICHAEL, Indiana University.

If the conjecture stated in this problem is true it is highly desirable to have a proof of it. It would be a significant contribution to the theory of prime numbers; for the theorem so obtained would afford a means of recursion by which a sequence of indefinitely increasing prime numbers could be written out consecutively. No such tool has yet been invented. Euler and Legendre sought in vain for analytical expressions which would serve just this purpose. Fermat believed, though he confessed that he was unable to prove, that he had found such an analytical expression in

$$2^{2n} + 1.$$

Euler pointed out the error of this opinion by showing that 641 is a factor of this number for the case $n = 5$.

From these historical facts one is led to suppose that the proof of the conjecture of this problem is probably difficult, in case the conjecture is true; if the conjecture is false, that would be shown by means of an example. The numbers to be dealt with, however, are so large that the construction of such an example (in case it exists) would perhaps be very tedious.

Note. A number of incorrect solutions of problem 198 have been received. Will our contributors please give it a more careful consideration? It is very probably not easy to solve.

EDITORS.

199. Proposed by R. P. LOCHNER, Philadelphia, Pa.

Find three integral squares, such that the sum of every two of them shall be a square.—Alsop's Algebra, 3d edition (Philadelphia, 1859), p. 296, Ex. 13.

SOLUTION BY ARTEMAS MARTIN, Washington, D. C.

Let x^2 , y^2 and z^2 be the required squares; then we must have

$$x^2 + y^2 = \square, \quad x^2 + z^2 = \square, \quad y^2 + z^2 = \square, \quad (1, 2, 3)$$

Assume $y = \frac{(m^2 - n^2)x}{2mn}$, $z = \frac{(p^2 - n^2)x}{2pn}$; then (1) and (2) are satisfied and

(3) becomes, after striking out common square factors,

$$p^2(m^2 - n^2)^2 + m^2(p^2 - n^2)^2 = \square, \quad (4)$$

which may be otherwise written

$$m^2p^2(m^2 + p^2) - 4m^2n^2p^2 + n^4(m^2 + p^2) = \square; \quad (5)$$

and (5) will be a square when

$$m^2 + p^2 = 4n^2. \quad (6)$$

Put $m = \frac{2n(r^2 - s^2)}{r^2 + s^2}$, $p = \frac{4nrs}{r^2 + s^2}$; then (6) is satisfied. Take $r = 2$, $s = 1$, and

we have $m = \frac{6n}{5}$, $p = \frac{8n}{5}$; $y = \frac{11x}{60}$, $z = \frac{39x}{80}$. Now take $x = 240$ and we have $y = 44$, $z = 117$, the least numbers known.